

# Impacts of Ship Emissions on Air Quality in Southern California

Satish Vutukuru, Joseph Ensberg, Donald Dabdub  
University of California Irvine

SECA Meeting  
December 11, 2007

---

---

---

---

---

---

---

## Outline

- Modeling Domain
- CIT Air Quality Model
- Ship Emissions
- Model Configuration
- Air Quality Impacts
- Conclusions, Next Steps

Computational Environmental Sciences Laboratory, UC Irvine

2/28

---

---

---


---

---

---

---

## Modeling Domain



Computational Grid: 80 x 30 horizontal cells, 5x5 km resolution, 5 vertical layers up to 1100 m

Computational Environmental Sciences Laboratory, UC Irvine

3/28

---

---

---

---

---

---

---


## CIT Model

- Initially developed at Cal Tech and is currently under continuous development at UCI
- Eularian photochemical grid model
- Incorporates state-of-science modules to model atmospheric processes
- The model is evaluated using data from several field campaigns
- The model is used for many applications to study air quality for the south coast and other regions

Computational Environmental Sciences Laboratory, UC Irvine

4/28

## Model Components



**Numerical Algorithms**  
QSTSE for advection  
PFISLM for aerosol dynamics

Computational Environmental Sciences Laboratory, UC Irvine

**CACM Mechanism**  
191 species: 120 fully integrated,  
67 pseudo-steady-state, and  
4 fixed concentration species

5/28

## CIT Model Evaluation

- CIT Model was validated using data from major field experiments.
  - Harley 1993 evaluated the performance for gas-phase species using data from 1987 SCAQS episode.
  - Meng 1998 compared aerosol predictions with observations from 1987 SCAQS episode.
  - Griffin 2002 compared organic aerosol predictions with 1993 field campaign observations.
  - Griffin 2006 applied model components (CACM, MPMPPO) to North East region and compared with observations

Computational Environmental Sciences Laboratory, UC Irvine

6/28

## CIT Model Applications


- CIT Model is extensively applied to South Coast region for air quality studies.
  - Effects of NO<sub>x</sub> and VOC controls on ambient particulate matter (*Nguyen, 2002*)
  - Modeling of sea-salt chemistry (*Knipping, 2002*)
  - Modeling of renoxification chemistry (*Knipping, 2002*)
  - Modeling the oxidative capacity (*Griffin, 2004*)
  - Calculation of incremental secondary organic aerosol reactivity (*Carreras-Sospedra, 2005*)
  - Distributed power generation studies (*Rodriguez, 2006*)
  - Secondary organic aerosol analysis (*Vutukuru, 2006*)

Computational Environmental Sciences Laboratory, UC Irvine

7/28

## Ship Emissions

- Spatially-resolved ship emissions for North America are obtained from Prof. Jim Corbett (U Delaware)
- Emissions are formatted to model resolution and size




Computational Environmental Sciences Laboratory, UC Irvine

8/28

## Ship Emissions


- Similarly, emissions of NO<sub>x</sub>, PM and TOG from ships for the base year 2002 are processed for a summer month
- Hourly emissions are calculated using uniform distribution of emissions throughout the day.



Computational Environmental Sciences Laboratory, UC Irvine

9/28

## Ship Emissions



Computational Environmental Sciences Laboratory, UC Irvine

10/28

---

---

---

---

---

---

---

---

## Ship Emission Speciation

- Emissions from for ROG and PM are speciated
- Chemical mechanisms in air quality models require emissions from individual species
- PM speciation also requires size distribution
- ARB speciation profiles are used to approximate speciation for ship emissions
  - combustion of distillate fuel

Computational Environmental Sciences Laboratory, UC Irvine

11/28

---

---

---

---

---

---

---

---

## Model Inputs

- Meteorological inputs are used from a 1987 SCAQS campaign.
  - A three-day episode that occurred from August 26 to August 29
  - A comprehensive dataset that is most representative of SoCAB's meteorological conditions
  - Extensively used for air quality studies in the region
- Baseline inventory for the year 1997 is scaled to 2002 levels using county-wide factors.
  - Baseline inventory was used in 2003 AQMP by SCAQMD

Computational Environmental Sciences Laboratory, UC Irvine

12/28

---

---

---

---

---

---


---

---

## Model Configuration

**O3: aq, Sf: ~ 0, Nt: ~ 0**

**O3: 40 ppb,  
Sf: ~ 0,  
Nt: ~ 0**



**O3: aq,  
Sf: ~ 0,  
Nt: ~ 0**

**O3: 40 ppb, Sf: ~ 0, Nt: ~ 0**

Computational Environmental Sciences Laboratory, UC Irvine

13/28

---

---

---

---


---

---

---

---

## Ambient Sulfates



**24-hour average concentration of ambient sulfates  
without ship emissions**

Computational Environmental Sciences Laboratory, UC Irvine

14/28

---

---

---

---


---

---

---

---

## Contribution from Ships



**Difference between 24-hour average concentration of  
ambient sulfates with and without ship emissions**

Computational Environmental Sciences Laboratory, UC Irvine

15/28

---

---

---

---


---

---

---

---

## Impact on Ambient Sulfates



Computational Environmental Sciences Laboratory, UC Irvine

16/28

---

---

---

---

---

---


---

---

---

---

## Contribution from Ships



Computational Environmental Sciences Laboratory, UC Irvine

17/28

---

---

---

---

---

---


---

---

---

---

## Impact on Ambient Nitrates



Computational Environmental Sciences Laboratory, UC Irvine

18/28

---

---

---

---

---

---


---

---

---

---

## Contribution from Ships



Difference between 24-hour average concentration of ambient nitrates with and without ship emissions

Computational Environmental Sciences Laboratory, UC Irvine

19/28


## Role of NO<sub>x</sub>

- NO<sub>x</sub> produces ozone which in turn produces OH radicals
- OH radicals and ozone play a key role in secondary particulate formation
- Relationship between NO<sub>x</sub> emissions and secondary particulate formation is highly non-linear

Computational Environmental Sciences Laboratory, UC Irvine

20/28

## Impact on Peak Ozone




Difference between peak ozone concentrations

Computational Environmental Sciences Laboratory, UC Irvine

21/28

## Role of NOx



Source: Khoi, 2002

**A Sensitivity Experiment:  
What is the impact we remove all NOx from ships?**

Computational Environmental Sciences Laboratory, UC Irvine

22/28

---

---

---

---


---

---

---

---

## Role of NOx



**Difference between peak ozone concentrations with zero Nox from ships**

Computational Environmental Sciences Laboratory, UC Irvine

23/28

---

---

---

---


---

---

---

---

## Role of NOx



**24-hour average concentration of ambient nitrates  
with zero NOx from ship emissions**

Computational Environmental Sciences Laboratory, UC Irvine

24/28

---

---

---

---

---


---

---

---



## Role of NOx



Computational Environmental Sciences Laboratory, UC Irvine

25/28

## Conclusions

- Impact of ship emissions
  - Up to  $1.5 \mu\text{g}/\text{m}^3$  for sulfates and  $12 \mu\text{g}/\text{m}^3$  for nitrates
  - Percentage contribution as high as 45% (Simi Valley area) for ambient sulfates and extends beyond Riverside (15% contribution)
  - Most impact is predicted at locations downwind of urban coastal areas
- NOx from ship emissions contribute to ozone significantly and secondary particulate formation

Computational Environmental Sciences Laboratory, UC Irvine

26/28

## Next Steps

- Refine assumptions based on discussions at this meeting
- Conduct sensitivity analyses
- Possibly conduct simulations for a future year.

Computational Environmental Sciences Laboratory, UC Irvine

27/28

## Acknowledgements

- Dongmin Luo and Ajith Kaduwela  
(California Air Resources Board)
- Prof. Jim Corbett (U Delaware)
- Tony Soeller, Joseph Farran  
(UCI Academic Computing Services)

Computational Environmental Sciences Laboratory, UC Irvine

28/28